

# Master equation

Koji Usami\*

(Dated: December 05, 2016)

We have learned how to treat the dissipative element, that is, the *environment*, quantum mechanically. Now we turn our attention to the *system* and learn how the dynamics of the system is affected by the environment. We shall start by considering, from the viewpoint of Schrödinger, a harmonic oscillator as the system, which is coupled to an Ohmic environment, and analyze the resultant master equation for the damped harmonic oscillator.

## I. "COARSE-GRAINED" TIME EVOLUTION [1]

Let us reexamine the LCR circuit from the viewpoint of Hamiltonian formalism. We shall assume the total Hamiltonian to be

$$H = H_s + H_b + H_I, \quad (1)$$

where  $H_s$ ,  $H_b$ , and  $H_I$  are the Hamiltonians of the LC circuit (the system), the transmission line (the Ohmic environment), and their interaction, and are respectively given by

$$H_s = \hbar\omega_0 \hat{a}^\dagger \hat{a} \quad (2)$$

$$H_b = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \hbar\omega \hat{c}^\dagger(\omega) \hat{c}(\omega) \quad (3)$$

$$\begin{aligned} H_I &= -i\hbar \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} (f(\omega) \hat{a}^\dagger \hat{c}(\omega) - f^*(\omega) \hat{a} \hat{c}^\dagger(\omega)) \\ &= -i\hbar \left[ \hat{a}^\dagger \underbrace{\int_{-\infty}^{\infty} \frac{d\omega}{2\pi} f(\omega) \hat{c}(\omega)}_{R^-} - \hat{a} \underbrace{\int_{-\infty}^{\infty} \frac{d\omega}{2\pi} f^*(\omega) \hat{c}^\dagger(\omega)}_{R^+} \right] \\ &= -i\hbar [\hat{a}^\dagger R^- - \hat{a} R^+]. \end{aligned} \quad (4)$$

Here the coupling strength  $f(\omega)$  has the dimension of  $\sqrt{\text{Angular frequency}}$  and will be identified to the familiar quantity later on.

Let us analyze the coupled system by focusing on the evolution of the density operator  $\rho$  of global system. The time evolution equation in the Schrödinger representation is

$$\frac{d}{dt} \rho(t) = \frac{1}{i\hbar} [H, \rho(t)]. \quad (5)$$

In the interaction representation with respect to  $H_s + H_b$ , it becomes

$$\frac{d}{dt} \tilde{\rho}(t) = \frac{1}{i\hbar} [\tilde{H}_I(t), \tilde{\rho}(t)], \quad (6)$$

where

$$\tilde{\rho}(t) = e^{i\frac{H_s+H_b}{\hbar}t} \rho(t) e^{-i\frac{H_s+H_b}{\hbar}t} \quad (7)$$

$$\begin{aligned} \tilde{H}_I(t) &= e^{i\frac{H_s+H_b}{\hbar}t} H_I e^{-i\frac{H_s+H_b}{\hbar}t} \\ &= -i\hbar \left[ \hat{a}^\dagger e^{i\omega_0 t} \tilde{R}^-(t) - \hat{a} e^{-i\omega_0 t} \tilde{R}^+(t) \right], \end{aligned} \quad (8)$$

---

\*Electronic address: [usami@qc.rcast.u-tokyo.ac.jp](mailto:usami@qc.rcast.u-tokyo.ac.jp)

where

$$\tilde{R}^-(t) = e^{i\frac{H_s+H_b}{\hbar}t} R_- e^{-i\frac{H_s+H_b}{\hbar}t} = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} f(\omega) \hat{c}(\omega) e^{-i\omega t} \quad (9)$$

$$\tilde{R}^+(t) = e^{i\frac{H_s+H_b}{\hbar}t} R_+ e^{-i\frac{H_s+H_b}{\hbar}t} = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} f^*(\omega) \hat{c}^\dagger(\omega) e^{i\omega t}. \quad (10)$$

The benefit of turning to the interaction representation is that if the interaction  $H_I$  is sufficiently small  $\tilde{\rho}(t)$  evolves slowly.

By formally integrating Eq. (6) we have

$$\tilde{\rho}(t + \Delta t) = \tilde{\rho}(t) + \frac{1}{i\hbar} \int_t^{t+\Delta t} dt' [\tilde{H}_I(t'), \tilde{\rho}(t')] \quad (11)$$

which leads to the following iterative relations:

$$\begin{aligned} \Delta\tilde{\rho}(t) &\equiv \tilde{\rho}(t + \Delta t) - \tilde{\rho}(t) \\ &= \frac{1}{i\hbar} \int_t^{t+\Delta t} dt' [\tilde{H}_I(t'), \tilde{\rho}(t')] \\ &= \frac{1}{i\hbar} \int_t^{t+\Delta t} dt' [\tilde{H}_I(t'), \tilde{\rho}(t)] + \left(\frac{1}{i\hbar}\right)^2 \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' [\tilde{H}_I(t'), [\tilde{H}_I(t''), \tilde{\rho}(t'')]] \\ &= \dots \end{aligned} \quad (12)$$

Let us now assume that the interaction between the system and the environment is small and retain the terms in Eq. (12) up to second order in  $\tilde{H}_I$ :

$$\Delta\tilde{\rho}(t) = \frac{1}{i\hbar} \int_t^{t+\Delta t} dt' [\tilde{H}_I(t'), \tilde{\rho}(t)] + \left(\frac{1}{i\hbar}\right)^2 \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' [\tilde{H}_I(t'), [\tilde{H}_I(t''), \tilde{\rho}(t)']]. \quad (13)$$

Now we make a crucial assumption. By writing the reduced density operators  $\tilde{\sigma}(t) = \text{Tr}_b \{\rho(t)\}$  and  $\tilde{\sigma}_b(t) = \text{Tr}_s \{\rho(t)\}$  for the system (LC) and environment (R), respectively, let us assume

$$\tilde{\rho}(t) = \tilde{\sigma}(t) \otimes \tilde{\sigma}_b(t), \quad (14)$$

meaning that the correlations between the system (LC) and the environment (R) at time  $t$  are neglected as they disappear very quickly compared with the typical time scale  $\Delta t$  we are interested in. Further assume that the environment is not affected by the coupling to the system and thus the environment is in a stationary state, that is,

$$\tilde{\sigma}_b(t) = \tilde{\sigma}_b(0) = \sigma_b \quad (15)$$

and thus

$$[\sigma_b, H_b] = 0. \quad (16)$$

Now, by tracing out the environmental degrees of freedom we have

$$\text{Tr}_b \{\Delta\tilde{\rho}(t)\} = \left(\frac{1}{i\hbar}\right)^2 \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' \text{Tr}_b \left\{ [\tilde{H}_I(t'), [\tilde{H}_I(t''), \tilde{\rho}(t)]] \right\}, \quad (17)$$

where

$$\text{Tr}_b \{\dots\} = \sum_n \langle n | \dots | n \rangle \quad (18)$$

with  $\{|n\rangle\}$  being the complete set of states for the environment. Here the first term in Eq. (13) disappears since

$$\begin{aligned} \text{Tr}_b \left\{ [\tilde{H}_I(t'), \tilde{\rho}(t)] \right\} &= \text{Tr}_b \left\{ [\tilde{H}_I(t'), \tilde{\sigma}(t) \otimes \sigma_b] \right\} \\ &= -i\hbar \left[ \hat{a}^\dagger e^{i\omega_0 t'} \text{Tr}_b \left\{ \tilde{R}^-(t') \sigma_b \right\} - \hat{a} e^{-i\omega_0 t'} \text{Tr}_b \left\{ \tilde{R}^+(t') \sigma_b \right\} \right] \tilde{\sigma}(t) \\ &\quad - i\hbar \tilde{\sigma}(t) \left[ \hat{a}^\dagger e^{i\omega_0 t'} \text{Tr}_b \left\{ \tilde{R}^-(t') \sigma_b \right\} - \hat{a} e^{-i\omega_0 t'} \text{Tr}_b \left\{ \tilde{R}^+(t') \sigma_b \right\} \right]. \end{aligned} \quad (19)$$

In case where the environment is in the thermal equilibrium

$$\mathrm{Tr}_b \left\{ \tilde{R}^+(t') \sigma_b \right\} = \mathrm{Tr}_b \left\{ \tilde{R}^-(t') \sigma_b \right\} = 0, \quad (20)$$

thus

$$\mathrm{Tr}_b \left\{ \left[ \tilde{H}_I(t'), \tilde{\rho}(t) \right] \right\} = 0. \quad (21)$$

The *coarse-grained* time evolution of  $\tilde{\sigma}(t)$  can then be given by

$$\begin{aligned} \frac{\Delta \tilde{\sigma}(t)}{\Delta t} &\equiv \frac{\mathrm{Tr}_b \{ \Delta \tilde{\rho}(t) \}}{\Delta t} \\ &= -\frac{1}{\hbar^2} \frac{1}{\Delta t} \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' \mathrm{Tr}_b \left\{ \left[ \tilde{H}_I(t'), \left[ \tilde{H}_I(t''), \tilde{\sigma}(t) \otimes \sigma_b \right] \right] \right\} \\ &= -\frac{1}{\hbar^2} \frac{1}{\Delta t} \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' \left[ \mathrm{Tr}_b \left\{ \tilde{H}_I(t') \tilde{H}_I(t'') (\tilde{\sigma}(t) \otimes \sigma_b) \right\} \right. \\ &\quad \left. - \mathrm{Tr}_b \left\{ \tilde{H}_I(t') (\tilde{\sigma}(t) \otimes \sigma_b) \tilde{H}_I(t'') \right\} \right. \\ &\quad \left. - \mathrm{Tr}_b \left\{ \tilde{H}_I(t'') (\tilde{\sigma}(t) \otimes \sigma_b) \tilde{H}_I(t') \right\} \right. \\ &\quad \left. + \mathrm{Tr}_b \left\{ (\tilde{\sigma}(t) \otimes \sigma_b) \tilde{H}_I(t'') \tilde{H}_I(t') \right\} \right]. \end{aligned} \quad (22)$$

## II. LINDBLAD FORM [1–3]

Let us further investigate the coarse-grained time evolution of the reduced density operator  $\sigma(t)$  in Eq. (22) to obtain the so-called *Lindblad form*, the non-unitary time evolution which nonetheless preserves complete positivity and trace of  $\sigma(t)$ . By plugging Eq. (8) into Eq. (22) we get

$$\begin{aligned} \frac{\Delta \tilde{\sigma}(t)}{\Delta t} &= \frac{1}{\Delta t} \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' [ \\ &\quad \mathrm{Tr}_b \left\{ \left( \tilde{a}^\dagger(t') \tilde{R}^-(t') - \tilde{a}(t') \tilde{R}^+(t') \right) \left( \tilde{a}^\dagger(t'') \tilde{R}^-(t'') - \tilde{a}(t'') \tilde{R}^+(t'') \right) (\tilde{\sigma}(t) \otimes \sigma_b) \right\} \\ &\quad - \mathrm{Tr}_b \left\{ \left( \tilde{a}^\dagger(t') \tilde{R}^-(t') - \tilde{a}(t') \tilde{R}^+(t') \right) (\tilde{\sigma}(t) \otimes \sigma_b) \left( \tilde{a}^\dagger(t'') \tilde{R}^-(t'') - \tilde{a}(t'') \tilde{R}^+(t'') \right) \right\} \\ &\quad - \mathrm{Tr}_b \left\{ \left( \tilde{a}^\dagger(t'') \tilde{R}^-(t'') - \tilde{a}(t'') \tilde{R}^+(t'') \right) (\tilde{\sigma}(t) \otimes \sigma_b) \left( \tilde{a}^\dagger(t') \tilde{R}^-(t') - \tilde{a}(t') \tilde{R}^+(t') \right) \right\} \\ &\quad + \mathrm{Tr}_b \left\{ (\tilde{\sigma}(t) \otimes \sigma_b) \left( \tilde{a}^\dagger(t'') \tilde{R}^-(t'') - \tilde{a}(t'') \tilde{R}^+(t'') \right) \left( \tilde{a}^\dagger(t') \tilde{R}^-(t') - \tilde{a}(t') \tilde{R}^+(t') \right) \right\} \\ &\quad ] \\ &= \frac{1}{\Delta t} \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' [ \\ &\quad - \langle \tilde{R}^-(t') \tilde{R}^+(t'') \rangle \tilde{a}^\dagger(t') \tilde{a}(t'') \tilde{\sigma}(t) - \langle \tilde{R}^+(t') \tilde{R}^-(t'') \rangle \tilde{a}(t') \tilde{a}^\dagger(t'') \tilde{\sigma}(t) \\ &\quad + \langle \tilde{R}^-(t'') \tilde{R}^+(t') \rangle \tilde{a}(t') \tilde{\sigma}(t) \tilde{a}^\dagger(t'') + \langle \tilde{R}^+(t'') \tilde{R}^-(t') \rangle \tilde{a}^\dagger(t') \tilde{\sigma}(t) \tilde{a}(t'') \\ &\quad + \langle \tilde{R}^-(t') \tilde{R}^+(t'') \rangle \tilde{a}(t'') \tilde{\sigma}(t) \tilde{a}^\dagger(t') + \langle \tilde{R}^+(t') \tilde{R}^-(t'') \rangle \tilde{a}^\dagger(t'') \tilde{\sigma}(t) \tilde{a}(t') \\ &\quad - \langle \tilde{R}^-(t'') \tilde{R}^+(t') \rangle \tilde{\sigma}(t) \tilde{a}^\dagger(t'') \tilde{a}(t') - \langle \tilde{R}^+(t'') \tilde{R}^-(t') \rangle \tilde{\sigma}(t) \tilde{a}(t'') \tilde{a}^\dagger(t') \\ &\quad ], \end{aligned} \quad (23)$$

where

$$\tilde{a}(t) = \hat{a} e^{-i\omega_0 t} \quad (24)$$

$$\tilde{a}^\dagger(t) = \hat{a}^\dagger e^{i\omega_0 t} \quad (25)$$

and

$$\langle F \rangle = \mathrm{Tr}_b \{ F \sigma_b \}, \quad (26)$$

for any operator  $F$  for the environment. Here the terms involving the products  $\langle \tilde{R}^+(t_1)\tilde{R}^+(t_2) \rangle$  and  $\langle \tilde{R}^-(t_1)\tilde{R}^-(t_2) \rangle$  are omitted for those produce zero when evaluating the expectation values with respect to the thermal state  $\sigma_b$ . By putting the reduced density operator  $\tilde{\sigma}(t)$  outside of the integral we have

$$\begin{aligned} \frac{\Delta\tilde{\sigma}(t)}{\Delta t} = & -\frac{1}{\Delta t} \left[ (\hat{a}^\dagger \hat{a} \tilde{\sigma}(t) - \hat{a} \tilde{\sigma}(t) \hat{a}^\dagger) \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' \left( \langle \tilde{R}^-(t')\tilde{R}^+(t'') \rangle e^{i\omega_0(t'-t'')} \right) \right. \\ & \left. + (\tilde{\sigma}(t) \hat{a} \hat{a}^\dagger - \hat{a}^\dagger \tilde{\sigma}(t) \hat{a}) \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' \left( \langle \tilde{R}^+(t'')\tilde{R}^-(t') \rangle e^{i\omega_0(t'-t'')} \right) \right] \\ & + h.c. \end{aligned} \quad (27)$$

We now calculate the two-time averages in Eq. (27). Note that from Eqs. (9) and (10) we have

$$\begin{aligned} \langle \tilde{R}^-(t')\tilde{R}^+(t'') \rangle &= \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \int_{-\infty}^{\infty} \frac{d\omega'}{2\pi} f(\omega) f^*(\omega') \underbrace{\langle \hat{c}(\omega)\hat{c}^\dagger(\omega') \rangle}_{\langle (n+1)2\pi\delta(\omega-\omega') \rangle} e^{-i\omega t'} e^{i\omega' t''} \\ &= \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} |f(\omega)|^2 \langle (n(\omega) + 1) \rangle e^{-i\omega(t'-t'')} \end{aligned} \quad (28)$$

and similarly

$$\begin{aligned} \langle \tilde{R}^+(t'')\tilde{R}^-(t') \rangle &= \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \int_{-\infty}^{\infty} \frac{d\omega'}{2\pi} f^*(\omega) f(\omega') \underbrace{\langle \hat{c}^\dagger(\omega)\hat{c}(\omega') \rangle}_{\langle n2\pi\delta(\omega-\omega') \rangle} e^{i\omega' t''} e^{-i\omega t'} \\ &= \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} |f(\omega)|^2 \langle n(\omega) \rangle e^{-i\omega(t'-t'')} \end{aligned} \quad (29)$$

where we used

$$[\hat{c}(\omega), \hat{c}^\dagger(\omega')] = 2\pi\delta(\omega - \omega') \quad (30)$$

and

$$\langle \hat{c}^\dagger(\omega)\hat{c}(\omega') \rangle = \text{Tr}_b \{ \hat{c}^\dagger(\omega)\hat{c}(\omega')\sigma_b \} = 2\pi\delta(\omega - \omega') \langle n(\omega) \rangle. \quad (31)$$

Since the two-time averages of  $\langle \tilde{R}^-(t')\tilde{R}^+(t'') \rangle$  and  $\langle \tilde{R}^+(t'')\tilde{R}^-(t') \rangle$  depend only on the time difference  $\tau \equiv t' - t''$  we can modify the domain of integration in Eq. (27) as shown as the shaded area in Fig. 1, that is,

$$\int_t^{t+\Delta t} dt' \int_t^{t'} dt'' \rightarrow \int_0^{\Delta t} d\tau \int_{t+\tau}^{t+\Delta t} dt'. \quad (32)$$




FIG. 1: Domain of integration: Eq. (32).




FIG. 2: Domain of integration: Eq. (33).

Now let us further assume that  $\langle \tilde{R}^-(t') \tilde{R}^+(t' - \tau) \rangle$  and  $\langle \tilde{R}^+(t' - \tau) \tilde{R}^-(t') \rangle$  are only non-zero within the narrow region as shown as the red-shaded area in Fig. 1 where  $\tau < \tau_c \ll \Delta t$ . This allows us to extend the domain of integration as shown as the shaded area in Fig. 2, i.e.,

$$\int_0^{\Delta t} d\tau \int_{t+\tau}^{t+\Delta t} dt' \rightarrow \int_0^{\infty} d\tau \int_t^{t+\Delta t} dt' \quad (33)$$

to give

$$\begin{aligned} \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' \left( \langle \tilde{R}^-(t') \tilde{R}^+(t'') \rangle e^{i\omega_0(t'-t'')} \right) &= \Delta t \int_0^{\infty} d\tau \left( \langle \tilde{R}^-(\tau) \tilde{R}^+(0) \rangle e^{i\omega_0\tau} \right) \\ &= \Delta t \int_0^{\infty} d\tau \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} |f(\omega)|^2 (\langle n(\omega) \rangle + 1) e^{-i(\omega - \omega_0)\tau} \end{aligned} \quad (34)$$

and similarly

$$\begin{aligned} \int_t^{t+\Delta t} dt' \int_t^{t'} dt'' \left( \langle \tilde{R}^+(t'') \tilde{R}^-(t') \rangle e^{i\omega_0(t'-t'')} \right) &= \Delta t \int_0^{\infty} d\tau \left( \langle \tilde{R}^+(0) \tilde{R}^-(\tau) \rangle e^{i\omega_0\tau} \right) \\ &= \Delta t \int_0^{\infty} d\tau \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} |f(\omega)|^2 \langle n(\omega) \rangle e^{-i(\omega - \omega_0)\tau}. \end{aligned} \quad (35)$$

These integrals seem to diverge at  $\omega = \omega_0$ . To prevent the divergence let us insert the convergence factor  $\epsilon$  into

Eqs. (34) and (35) and perform integrations

$$\begin{aligned}
& \lim_{\epsilon \rightarrow 0^+} \int_0^\infty d\tau \int_{-\infty}^\infty \frac{d\omega}{2\pi} |f(\omega)|^2 (\langle n(\omega) \rangle + 1) e^{i[(\omega_0 - \omega) + i\epsilon]\tau} \\
&= i \int_{-\infty}^\infty \frac{d\omega}{2\pi} |f(\omega)|^2 (\langle n(\omega) \rangle + 1) \left( -i \lim_{\epsilon \rightarrow 0^+} \int_0^\infty d\tau e^{i[(\omega_0 - \omega) + i\epsilon]\tau} \right) \\
&= i \int_{-\infty}^\infty \frac{d\omega}{2\pi} |f(\omega)|^2 (\langle n(\omega) \rangle + 1) \underbrace{\left( \lim_{\epsilon \rightarrow 0^+} \frac{1}{(\omega_0 - \omega) + i\epsilon} \right)}_{G_+(\omega_0)}
\end{aligned} \tag{36}$$

$$\begin{aligned}
& \lim_{\epsilon \rightarrow 0^+} \int_0^\infty d\tau \int_{-\infty}^\infty \frac{d\omega}{2\pi} |f(\omega)|^2 \langle n(\omega) \rangle e^{i[(\omega - \omega_0) + i\epsilon]\tau} \\
&= i \int_{-\infty}^\infty \frac{d\omega}{2\pi} |f(\omega)|^2 \langle n(\omega) \rangle \left( i \lim_{\epsilon \rightarrow 0^+} \int_0^\infty d\tau e^{i[(\omega - \omega_0) + i\epsilon]\tau} \right) \\
&= i \int_{-\infty}^\infty \frac{d\omega}{2\pi} |f(\omega)|^2 \langle n(\omega) \rangle \underbrace{\left( \lim_{\epsilon \rightarrow 0^+} \frac{1}{(\omega_0 - \omega) + i\epsilon} \right)}_{G_+(\omega_0)}.
\end{aligned} \tag{37}$$

We now invoke the so-called *Dirac identity*,

$$G_+(\omega_0) \equiv \lim_{\epsilon \rightarrow 0^+} \frac{1}{(\omega_0 - \omega) + i\epsilon} = \frac{\mathcal{P}}{\omega_0 - \omega} - i\pi\delta(\omega_0 - \omega) \tag{38}$$

to get

$$\begin{aligned}
i \int_{-\infty}^\infty \frac{d\omega}{2\pi} |f(\omega)|^2 (\langle n(\omega) \rangle + 1) G_+(\omega_0) &= i \int_{-\infty}^\infty \frac{d\omega}{2\pi} |f(\omega)|^2 (\langle n(\omega) \rangle + 1) \left( \frac{\mathcal{P}}{\omega_0 - \omega} - i\pi\delta(\omega_0 - \omega) \right) \\
&= i \underbrace{\mathcal{P} \int_{-\infty}^\infty \frac{d\omega}{2\pi} \frac{|f(\omega)|^2 (\langle n(\omega) \rangle + 1)}{\omega_0 - \omega}}_{\Delta' + \Delta} \\
&\quad + \frac{1}{2} \underbrace{\int_{-\infty}^\infty d\omega |f(\omega)|^2 (\langle n(\omega) \rangle + 1) \delta(\omega_0 - \omega)}_{\Gamma' + \Gamma} \\
&= i(\Delta' + \Delta) + \frac{\Gamma' + \Gamma}{2}
\end{aligned} \tag{39}$$

$$\begin{aligned}
i \int_{-\infty}^\infty \frac{d\omega}{2\pi} |f(\omega)|^2 \langle n(\omega) \rangle G_+(\omega_0) &= i \int_{-\infty}^\infty \frac{d\omega}{2\pi} |f(\omega)|^2 \langle n(\omega) \rangle \left( \frac{\mathcal{P}}{\omega_0 - \omega} - i\pi\delta(\omega - \omega_0) \right) \\
&= i \underbrace{\mathcal{P} \int_{-\infty}^\infty \frac{d\omega}{2\pi} \frac{|f(\omega)|^2 \langle n(\omega) \rangle}{\omega - \omega_0}}_{\Delta'} \\
&\quad + \frac{1}{2} \underbrace{\int_{-\infty}^\infty d\omega |f(\omega)|^2 \langle n(\omega) \rangle \delta(\omega - \omega_0)}_{\Gamma'} \\
&= i\Delta' + \frac{\Gamma'}{2},
\end{aligned} \tag{40}$$

where  $\mathcal{P}$  is the principal part integral. For the function  $F(x)$  which diverges at the point  $x = x_0$ , it is defined by

$$\mathcal{P} \int_a^b F(x) dx = \lim_{\delta \rightarrow 0} \left( \int_a^{x_0 - \delta} F(x) dx + \int_{x_0 + \delta}^b F(x) dx \right). \tag{41}$$

By plugging these expressions in Eq. (27) we have

$$\begin{aligned}
\frac{\partial \tilde{\sigma}(t)}{\partial t} &= - \left[ (\hat{a}^\dagger \hat{a} \tilde{\sigma}(t) - \hat{a} \tilde{\sigma}(t) \hat{a}^\dagger) \left( i(\Delta' + \Delta) + \frac{\Gamma' + \Gamma}{2} \right) \right. \\
&\quad + (\tilde{\sigma}(t) \hat{a} \hat{a}^\dagger - \hat{a}^\dagger \tilde{\sigma}(t) \hat{a}) \left( i\Delta' + \frac{\Gamma'}{2} \right) \\
&\quad + (\tilde{\sigma}(t) \hat{a}^\dagger \hat{a} - \hat{a} \tilde{\sigma}(t) \hat{a}^\dagger) \left( -i(\Delta' + \Delta) + \frac{\Gamma' + \Gamma}{2} \right) \\
&\quad \left. + (\hat{a} \hat{a}^\dagger \tilde{\sigma}(t) - \hat{a}^\dagger \tilde{\sigma}(t) \hat{a}) \left( -i\Delta' + \frac{\Gamma'}{2} \right) \right] \\
&= -\frac{i}{\hbar} [(\hbar \Delta \hat{a}^\dagger \hat{a}), \tilde{\sigma}(t)] + \frac{\Gamma' + \Gamma}{2} (2\hat{a} \tilde{\sigma}(t) \hat{a}^\dagger - \hat{a}^\dagger \hat{a} \tilde{\sigma}(t) - \tilde{\sigma}(t) \hat{a}^\dagger \hat{a}) \\
&\quad + \frac{\Gamma'}{2} (2\hat{a}^\dagger \tilde{\sigma}(t) \hat{a} - \hat{a} \hat{a}^\dagger \tilde{\sigma}(t) - \tilde{\sigma}(t) \hat{a} \hat{a}^\dagger). \tag{42}
\end{aligned}$$

Returning to the Schödinger representation we arrive at the following *master equation*:

$$\begin{aligned}
\frac{\partial \sigma(t)}{\partial t} &= -\frac{i}{\hbar} [\hbar(\omega_0 + \Delta) \hat{a}^\dagger \hat{a}, \sigma(t)] + \frac{\Gamma' + \Gamma}{2} (2\hat{a} \sigma(t) \hat{a}^\dagger - \hat{a}^\dagger \hat{a} \sigma(t) - \sigma(t) \hat{a}^\dagger \hat{a}) \\
&\quad + \frac{\Gamma'}{2} (2\hat{a}^\dagger \sigma(t) \hat{a} - \hat{a} \hat{a}^\dagger \sigma(t) - \sigma(t) \hat{a} \hat{a}^\dagger). \tag{43}
\end{aligned}$$

We can now indentify  $\Gamma$  defined in Eqs. (39) and (40) by

$$\Gamma = \int_{-\infty}^{\infty} d\omega |f(\omega)|^2 \delta(\omega - \omega_0) \sim |f(\omega_0)|^2 \tag{44}$$

as the Einstein-*A*-coefficient-like spontaneous emission rate for the LC circuit. On the other hand  $\Gamma'$  defined in Eqs. (39) and (40) by

$$\Gamma' = \int_{-\infty}^{\infty} d\omega |f(\omega)|^2 \langle n(\omega) \rangle \delta(\omega - \omega_0) \sim |f(\omega_0)|^2 \langle n(\omega_0) \rangle = \langle n(\omega_0) \rangle \Gamma \tag{45}$$

is identified as the Einstein-*B*-coefficient-like stimulated emission rate (and also absorption rate).

It also becomes clear that  $\Delta$  appeared in Eqs. (39) and (40) is the Lamb-shift-like *spontaneous* radiative shift occurring without any photons in the environment, and  $\Delta'$  is the radiative shift due to real photons in the environment, which in fact *disappears* in Eq. (43) in a course of calculation (see Eq. (42)).

The master equation, Eq. (43), assumes the *Lindblad form*, which, in general, is written as

$$\begin{aligned}
\frac{\partial \sigma(t)}{\partial t} &= -\frac{i}{\hbar} [\mathcal{H}, \sigma(t)] + \frac{1}{2} \sum_J \left( 2A_J \sigma(t) A_J^\dagger - A_J^\dagger A_J \sigma(t) - \sigma(t) A_J^\dagger A_J \right) \\
&= -\frac{i}{\hbar} [\mathcal{H}, \sigma(t)] + \frac{1}{2} \sum_J \mathcal{L}_D[A_J] \sigma(t), \tag{46}
\end{aligned}$$

with  $\mathcal{H}$  being some hermitial operator and  $A_J$  being any operators (called *collapse operators*), which are responsible for the irreversible non-unitary evolution of  $\sigma(t)$ . The Lindblad form, however, assures to preserve the complete positivity and trace of the reduced density operator  $\sigma(t)$ . Here the *Lindblad superoperator*  $\mathcal{L}_D[A_J] \sigma(t)$  is defined by

$$\mathcal{L}_D[A_J] \sigma(t) = 2A_J \sigma(t) A_J^\dagger - A_J^\dagger A_J \sigma(t) - \sigma(t) A_J^\dagger A_J. \tag{47}$$

The master equation (43) is then written with the Lindblad superoperators as

$$\frac{\partial \sigma(t)}{\partial t} = -\frac{i}{\hbar} [\hbar(\omega_0 + \Delta) \hat{a}^\dagger \hat{a}, \sigma(t)] + \frac{1}{2} \mathcal{L}_D[\sqrt{\Gamma' + \Gamma} \hat{a}] \sigma(t) + \frac{1}{2} \mathcal{L}_D[\sqrt{\Gamma'} \hat{a}^\dagger] \sigma(t). \tag{48}$$




FIG. 3: Physical interpretations of 4 terms in Eq. (49).

### III. RATE EQUATION [1]

Now we shall see more explicitly that  $\Gamma = |f(\omega_0)|^2$  in Eq. (44) is related to the Einstein-A-coefficient-like spontaneous emission rate. Let us first evaluate the master equation (43) in the Fock-state bases  $\{|N\rangle\}$  for the system:

$$\begin{aligned} \frac{dP(N, t)}{dt} &= \frac{\Gamma + \Gamma'}{2} (2\langle N|\hat{a}\sigma(t)\hat{a}^\dagger|N\rangle - \langle N|\hat{a}^\dagger\hat{a}\sigma(t)|N\rangle - \langle N|\sigma(t)\hat{a}^\dagger\hat{a}|N\rangle) \\ &\quad + \frac{\Gamma'}{2} (2\langle N|\hat{a}^\dagger\sigma(t)\hat{a}|N\rangle - \langle N|\hat{a}\hat{a}^\dagger\sigma(t)|N\rangle - \langle N|\sigma(t)\hat{a}\hat{a}^\dagger|N\rangle) \\ &= \underbrace{\Gamma_\uparrow}_{\Gamma'} NP(N-1) - \underbrace{\Gamma_\uparrow}_{\Gamma'} (N+1)P(N) - \underbrace{\Gamma_\downarrow}_{\Gamma+\Gamma'} NP(N) + \underbrace{\Gamma_\downarrow}_{\Gamma+\Gamma'} (N+1)P(N+1), \end{aligned} \quad (49)$$

where we used the abbreviations  $P(N, t) = \langle N|\sigma(t)|N\rangle$ ,  $P(N-1, t) = \langle N-1|\sigma(t)|N-1\rangle$ , and  $P(N+1, t) = \langle N+1|\sigma(t)|N+1\rangle$ . Here the first term in the right hand side of Eq. (43) naturally disappears since it represents the conservative Hamiltonian and thus contributes nothing to the dynamics of Fock states (energy eigenstates). Consequently, this equation contains only the diagonal elements of the density operator and the effect of the radiative shifts  $\Delta$  and  $\Delta'$  are absent. There are 4 terms in Eq. (49), each of which has a clear physical meaning as depicted in Fig. 3. The first term represents the population gain from the  $(N-1)$ -photon Fock state due to the absorption at the rate of  $N\Gamma'$  (the lower blue arrow in Fig. 3). The second term represents the population loss from the  $N$ -photon Fock state due to the absorption at the rate of  $(N+1)\Gamma'$  (the upper blue arrow in Fig. 3). The third and the fourth terms represent the spontaneous and stimulated emissions from the  $N$ -photon Fock state into the  $(N-1)$ -photon Fock state at the rate of  $N(\Gamma'+\Gamma)$  (the lower red arrow in Fig. 3) and from the  $(N+1)$ -photon Fock state into the  $N$ -photon Fock state at the rate of  $(N+1)(\Gamma'+\Gamma)$  (the upper red arrow in Fig. 3), respectively.

We have then the following rate equation:

$$\begin{aligned} \frac{d}{dt}\langle N \rangle &\equiv \sum_N N \frac{dP(N, t)}{dt} \\ &= \Gamma_\uparrow \sum_N (N+1)P(N) - \Gamma_\downarrow \sum_N NP(N), \\ &= \Gamma_\uparrow \langle N+1 \rangle - \Gamma_\downarrow \langle N \rangle \\ &= -(\Gamma_\downarrow - \Gamma_\uparrow) \langle N \rangle + \Gamma_\uparrow \\ &= -\Gamma \langle N \rangle + \Gamma' \\ &= -\Gamma \langle N \rangle + \Gamma \langle n(\omega_0) \rangle, \end{aligned} \quad (50)$$

where Eq. (45) is used in the last equation. Thus, as we anticipated,  $\Gamma$  can be considered as the spontaneous emission rate. Note that the bosonic enhancement factors are canceled to get the damping rate independent on the initial excitations. This cancellation stems from the fact that the harmonic oscillators have an equidistant energy-level structure. The steady state condition  $\frac{d}{dt}\langle N \rangle = 0$  gives us the steady-state occupation number of the LC photons:

$$\bar{N} = \frac{\Gamma'}{\Gamma} = \langle n(\omega_0) \rangle, \quad (51)$$

that is, the averaged photon number of the environment  $\langle n(\omega_0) \rangle$ .

The general solution of the rate equation Eq. (50) can be given by

$$\langle N(t) \rangle = \langle N(0) \rangle e^{-\Gamma t} + \langle n(\omega_0) \rangle (1 - e^{-\Gamma t}). \quad (52)$$

#### IV. PROBLEM

##### A. Fokker-Planck equation [1]

In Sec. III we evaluate the master equation (43) in the Fock-state bases  $\{|N\rangle\}$  to obtain the rate equation. When we evaluate the master equation (43) in the coherent-state bases  $\{|\alpha\rangle\}$ , we arrive at the *Fokker-Planck equation* for the quasi-probability density called *P-function*,  $P_N(\beta, \beta^*, t)$ . Here the P-function can be used to represent the density operator  $\sigma(t)$  as

$$\sigma(t) = \int d^2\beta P_N(\beta, \beta^*, t) |\beta\rangle\langle\beta|, \quad (53)$$

where the coherent-state bases  $\{|\beta\rangle\}$  are defined as

$$\hat{a}|\beta\rangle = \beta|\beta\rangle \quad (54)$$

$$\langle\beta|\hat{a}^\dagger = \beta^*\langle\beta|. \quad (55)$$

Deduce the Fokker-Planck equation

$$\begin{aligned} \frac{\partial}{\partial t} P_N(\beta, \beta^*, t) &= \left( \frac{\Gamma}{2} + i(\omega_0 + \Delta) \right) \frac{\partial}{\partial \beta} [\beta P_N(\beta, \beta^*, t)] \\ &+ \left( \frac{\Gamma}{2} - i(\omega_0 + \Delta) \right) \frac{\partial}{\partial \beta^*} [\beta^* P_N(\beta, \beta^*, t)] \\ &+ \Gamma' \frac{\partial^2}{\partial \beta \partial \beta^*} P_N(\beta, \beta^*, t). \end{aligned} \quad (56)$$

from Eq. (43) with Eq. (53).

Check that the following Gaussian function

$$P_N(\beta, \beta^*, t) = \frac{1}{\pi \langle n(\omega_0) \rangle (1 - e^{-\Gamma t})} \exp \left( - \frac{\left[ \beta - \beta_0 e^{-\left(\frac{\Gamma}{2} + i(\omega_0 + \Delta)t\right)} \right] \left[ \beta^* - \beta_0^* e^{-\left(\frac{\Gamma}{2} - i(\omega_0 + \Delta)t\right)} \right]}{\langle n(\omega_0) \rangle (1 - e^{-\Gamma t})} \right) \quad (57)$$

constitutes the solution of Eq. (56). Note that for  $t = 0$  it becomes  $\delta(\beta - \beta_0)\delta(\beta^* - \beta_0^*)$ . Show the function  $P_N(\beta, \beta^*, t)$  for  $t = \infty$ .

- 
- [1] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, *Atom-Photon Interactions* (John Wiley and Sons, New York, 1992).  
 [2] S. M. Barnett and P. M. Radmore, *Methods in Theoretical Quantum Optics* (Oxford University Press, New York, 1997).  
 [3] C. W. Gardiner and P. Zoller, *Quantum Noise*, 2nd ed. (Springer, Berlin, 2000).